

Patient No.	Age (years)	Sex	Tumor Location	Tumor Type	Size of Defect (cm ²)	Reconstructive Procedure	Follow-up (months)	Functional Score (%)
1	11	M	Medial aspect forefoot	Malignant Peripheral Nerve Sheath Tumor	(6 x 8 cm) 48 cm ²	Cross-leg flap (AWED)	91	100%
2	56	M	Lateral Heel	High Grade Liposarcoma	(26 x 7 cm) 182 cm ²	Rectus Abdominis Free Flap (AWED)	67	96.7%
3	21	F	Weight-bearing Heel	Clear Cell Sarcoma	(9 x 7 cm) 63 cm ²	Reverse Sural Artery Flap (DOD)	60	93.3%
4	40	F	Medial to Lateral Heel	Fibrosarcoma	(12 x 13 cm) 156 cm ²	Reverse Sural Artery Flap	56	90%
5	43	F	Medial Ankle	Synovial Sarcoma	(12 x 15 cm) 180 cm ²	Reverse Sural Artery Flap	17	87%
6	47	M	Medial Mid foot	Malignant Fibrous Histiocytoma	(9 x 13 cm) 117 cm ²	Reverse Sural Artery Flap	4	90%
Average	36.3 (11-56 years)				124.3 cm ²		49.2 (4-91 months)	92.8%

Table 2 Pertinent data on the six patients.